Spin polarization
and Einstein-Podolsky-Rosen paradox
in the MC event records

Z. Wąs
Institute of Nuclear Physics, Kraków, Poland

Main Topics:

- **Usual approach: helicity states.**
- **When it works?**
- **When it breaks?**
- **When it breaks badly?**
- **Why it does not matter in this case?**
- **Wen it may matter a lot, and constraints on the data structure for backup solution.**

Slides are (will be) on http://home.cern.ch/wasm

Helicity states.

For (nearly) all Monte Carlo architectures we use:

Stadard approach

1. Particles have properties.
2. Particles have origin.
3. Particles have decay products.

Used everywhere:

`HEPEVT, LUJETS, PYJETS, HEPEUP, HEPMC, ...`

but nonetheless ...

Let us have a critical look at it with the help of examples and with the special emphasis on breaking point and when the data structures organized around 'particles' can lead to the trap.

Stadard approach

Example: KORALZ at LEP, Monte Carlo program for production of τ-lepton pairs and decay, including QED bremssstrahlung corrections

S. Jadach, B.F.L. Ward, Z. Was Comput. Phys. Commun. 79 (1994) 503

- Generates first pair of τ leptons
- Generates helicity states of both τ^+ and τ^- i.e. approximation is used
- Provides helicty states and relation between τ's restframe and LAB to TAUOLA
- TAUOLA performs decay of 100 % polarized τ's.
- In (nearly) all cases at LEP such approach was sufficient.

That is standard solution used very often, advocated in general idea of `HEPEVT` common block since many years and also recently in discussions for e.g. data structure for LHC and c++.

Helicity states.

If something works in one place why not ...

Universal interface for TAUOLA –

library for τ lepton decay.

T. Pierzchala, E. Richter-Was, M. Worek, Z. Was Acta. Phys. Polon. B32 (2001) 1277.

- Search for τ's over `HEPEVT` common block filled by any MC program
- Look through the origin of τ lepton: $Z, \gamma, W, h, h^{\pm}$,

$e^+e^-(u\bar{u}, d\bar{d}) \to \tau^+\tau^-$

- Eventually calculate effective incoming beams
- Generate τ helicity states
- Generate τ decay(s)
- Complete event stored in `HEPEVT` common block with the decay products.

Helicity states.

τ-polarization in $e^+e^- \to \tau^+\tau^-$

Tests of the TAUOLA universal interface, preliminary results.
The τ lepton polarization as a function od $\cos\theta$ for the $e^+e^- \to \tau^+\tau^-$ process.



The function used in the interface is taken form KORALZ.

It looks after implementation (equally) OK.

Test at τ decay products level

Tests of the TAUOLA universal interface, preliminary results.
The π energy spectrum in the Z/γ rest frame. $\sqrt{s} \simeq M_Z$



1

$\pi^+ \pi^-$ Energy-Energy correlations. $\sqrt{s} \simeq M_Z$



2

To make spin effects large, we take sample of both τ' decaying into $\pi\nu$. Standard tests are getting reproduced qualitativelly.

We will continue with the more sophisticated tests in future. In particular we need to test the algorithm for matching bremsstrahlung.

Helicity states.

τ-polarization in $q\bar{q} \to \tau^+\tau^-$

Now let us use the same functions for quarks:

Tests of the TAUOLA universal interface, preliminary results
The τ lepton polarization as a function od $\cos\theta$ for the $u\bar{u} \to \tau^+\tau^-$ process.



2

Tests of the TAUOLA universal interface, preliminary results.
The τ lepton polarization as a function od $\cos\theta$ for the $d\bar{d} \to \tau^+\tau^-$ process.



3

KORALZ

It was based/on the careful downgrading from the full treatment of spin in KORALB (S. Jadach, Z. Was Comput. Phys. Commun. 64 (1991) 267) program for lower than LEP energies and first order radiative corrections only.
Tests of the validity of KORALZ spin treatments were ALWAYS available.

Anyway at LEP it was numerically important to have it only twice:

- For the measurement of transverse spin correlations and weak tests of the Bell inequalities
- For the measurement of the τ lepton lifetime using impact parameters difference method.

Universal interface of TAUOLA

Nothing like such possibility of tests exists,
except comparisons with KORALB KORALZ again ...

Anyway we feel confident

Should we always?

PHOTOS

E. Barberio, B. van Eijk, Z. Was, Comput. Phys. Commun. (1991) .

It was developped starting from the careful downgrading from the full treatment of M.E. in MUSTRAAL (F. Berends, R. Kleiss, S. Jadach, Comput. Phys. Commun. (1982)) For the decay of Z to obtain 'factorizable' form.

Extended to work for the decay of 'any' particle or resonance.

It heavily uses mother-daughter relations in `HEPEVT`

- Thus it is bound to be not better than LL (may be NLL)
- Safety backup: Comparisons with M.E. Monte Carlos whenever available.
- E.g. $\tau \to e\nu\bar{\nu}\gamma$, $\tau \to \pi\nu\gamma$, $Z \to \mu^+\mu^-\gamma(\gamma)$, $gg \to t\bar{t}\gamma\gamma$.

In many cases tests are not available ...,

DIRTY OBSERVABLE

Aim:

- Observable that produces unexpected peaks out of background expected to be flat.
- Starting point: $s\bar{s}$ state in $e^+e^- \rightarrow s\bar{s}c\bar{c}$ at 350 GeV ($c\bar{c}$ lost in the beam pipe).
- This is not realistic observable, but in principle realizable experimentally.
- Let us look at it with:

(i) full ME.,

(ii) CC03: WW prodction and decay

(iii) CC03 no transverse spin correlations.