

# Have we missed a Higgs at LEP?

**J. Alcaraz**  
CIEMAT/CERN



**ICHEP02**  
July 26, 2002

(Flavour independent searches and anomalous production and decay of Higgs bosons at LEP)


- ◆ Introduction. Dedicated searches
- ◆ Relaxing the assumptions
  - “All h,A” possibilities
  - “Flavour-independent” searches
  - “Model-independent” deviations
  - “Decay-independent” searches
- ◆ Summary

Yukawa searches for low mass Higgses

Triggered by





Higgs signature



$$4 \text{ GeV} < m_h, m_A < 12 \text{ GeV}$$

Large couplings to down-type quarks


$h, A \rightarrow b\bar{b}$  kinematically forbidden



Disfavours a low mass Higgs explanation to any large  $\delta a_\mu$  discrepancy


*Low  $m_A$ , high  $m_h$*

Initially triggered by




**Problematic region**

Higgs signature




$m_h$  large,  $m_A < 10$  GeV




Highly boosted A  $\Rightarrow$  1 jet-like decay

*OPAL: No excess observed*




Most of the accessible parameter space excluded

Can we do it better? Two Higgs Doublet models






## Upper bounds in terms of $\sigma/\sigma_{ref}$




95% CL excluded regions for  $\sigma/\sigma_{ref} = 1$






# $H \rightarrow b\bar{b}$ is really dominant?




- ◆ Main assumption: Higgs decays hadronically
- ◆ Main change in the studies: no b-tagging requirements

# "Flavour-independent" searches: Results




Similar sensitivities for all four LEP experiments

# LEP PRELIMINARY



$m_H > 113$  GeV at 95% CL

"Flavour-independent" searches: do not forget the  $\tau$




$m_H > 109.1 \text{ GeV}$  at 95% CL (ALEPH)  
(100% BR into  $q\bar{q}$  or  $\tau^+ \tau^-$ , SM production)

# Higgs deviations from the SM from $\Lambda \gg \sqrt{s}$

- ◆ If the SM is an approximately correct description of Nature, the “true theory” from a new physics scale  $\Lambda$  looks at low energies like (flavour violations are neglected):


$$\mathcal{L} = \mathcal{L}_{SM} + \sum_{n=6}^{\infty} \frac{1}{\Lambda^{n-4}} \sum_j \mathcal{O}_{nj}$$

- ◆ Already at dimension 6 some CP conserving anomalous HVV couplings, not excluded by LEP1 or lower energy experiments, may appear:



$$\begin{aligned} \mathcal{L} = \mathcal{L}_{SM} &+ g_{H\gamma\gamma} H A_{\mu\nu} A^{\mu\nu} + g_{HZ\gamma}^{(1)} A_{\mu\nu} Z^\mu \partial^\nu H + g_{HZ\gamma}^{(2)} H A_{\mu\nu} Z^{\mu\nu} \\ &+ g_{HZZ}^{(1)} Z_{\mu\nu} Z^\mu \partial^\nu H + g_{HZZ}^{(2)} H Z_{\mu\nu} Z^{\mu\nu} + g_{HZZ}^{(3)} H Z_\mu Z^\mu + \dots \end{aligned}$$


Effect: Renormalization of the Higgs field, do not change BRs  
 Search: reinterpretation of SM Higgs searches ( $\xi^2 = \sigma^{ZH} / \sigma_{SM}^{ZH}$ )





Anomalous Higgs couplings  $\Delta g_1^Z$ ,  $\Delta \kappa_\gamma$  (usual  $WWV$  couplings)

Effect: Change in  $HWW$ ,  $HZZ$ ,  $HZ\gamma$  couplings

Searches:  $e^+e^- \rightarrow HZ$  and  $e^+e^- \rightarrow H\gamma \rightarrow Z\gamma\gamma$




Anomalous Higgs couplings  $\Delta g_1^Z, \Delta \kappa_\gamma$ : **NO EXCESS**





$|\Delta g_1^Z|, |\Delta \kappa_\gamma| \lesssim 0.5$  at 95% CL,  $m_H \lesssim 160$  GeV



Effect: Change everything, but mostly  $H\gamma\gamma$  and  $HZ\gamma$  couplings  
 Searches:  $e^+e^- \rightarrow H\gamma$ ,  $\gamma\gamma \rightarrow H$ , ... , with  $H \rightarrow \gamma\gamma$  or  $H \rightarrow Z\gamma$




Anomalous Higgs couplings  $d$ ,  $d_B$ , NO EXCESS



$|d| \lesssim 0.1, |d_B| \lesssim 0.05$  at 95% CL,  $m_H \lesssim 160$  GeV


Alternative interpretation of  $d$ ,  $d_B$  limits



Limits to  $Br(H \rightarrow \gamma\gamma)$  and  $Br(H \rightarrow Z\gamma)$  in the region of LHC interest



“Decay-independent” searches

And if we do not want to assume anything . . .



- ◆ Aim: Efficiency should not depend on what happens on the Higgs side
- ◆ How: Use only  $Z \rightarrow e^+e^-$ ,  $Z \rightarrow \mu^+\mu^-$ , maybe  $Z \rightarrow \nu\bar{\nu}$

*OPAL*: No excess found





For  $\sigma/\sigma_{SM} = 1$ ,  $m_H > 81$  GeV at 95% CL

# And if the Higgs is too wide . . .

- ◆ Stealthy Higgs Model: Higgs singlets (phions) do not interact with matter
- ◆ But they can interact with the SM Higgs doublet:

$$\Delta\Gamma_H \rightarrow \frac{\omega^2 v^2}{32\pi m_H^2}, \quad N_{phions} \rightarrow \infty$$



For  $m_H \approx \Gamma_H$ ,  $m_H \gtrsim 50$  GeV at 95% CL

- ◆ Big efforts by the LEP Collaborations to be sure that we are not missing **ANY Higgs**
- ◆ Interesting studies in different directions by the different Collaborations
- ◆ For the moment, wherever we look for:  
**NO EVIDENCE FOR A FUNDAMENTAL SCALAR**