$F_{2,hadr}^{\gamma}$ and $\sigma_{\gamma^*\gamma^*}$

Results from L3 and Opal

Daniel Haas

University of Basel

April 2000

DIS 2000 – Liverpool

Overview

- $F_{2,hadr}^{\gamma}$ at L3 (Single Tag)
  - Available x – Q^2-range
  - Unfolding
  - Monte Carlo
  - Modeling Dependence
  - Results from L3 - Low and Medium Q^2
  - Results from L3 - High Q^2
- $\sigma_{\gamma^*\gamma^*}$ (Double-Tag)
  - Kinematics
  - BFKL
  - Results from L3
- Conclusions

F_2^γ : $x - Q^2$ at L3

Accessible Region

	$\sqrt{s}$ GeV	Q^2 GeV^2	$< P^2 >$ GeV^2	θ_{tag} mrad	x
L3 Low Q^2 (1)	91	1.2 - 9	0.08	26 - 66	0.002 - 0.2
L3 Medium Q^2 (2)	183	9 - 30	0.08	30 - 66	0.01 - 0.3
L3 High Q^2 Single Tag (3)	91	40 - 500	0.014	200 - 600	0.05 - 0.98
L3 High Q^2 Double Tag (3)	91	40 - 500	1 - 8	200 - 600	0.05 - 0.98

Used Sample and Cuts

	$\mathcal{L}$ pb^{-1}	Evts	Bkg	E_{tag} GeV	W_{vis} GeV
L3 Low Q^2	140	9928	355	> 35	3 - 40
L3 Medium Q^2	52	1694	71	> 70	> 3
L3 High Q^2 Single Tag	120	496	40	> 14	> 3
L3 High Q^2 Double Tag	120	43	< 2	> 14	> 3

(1) Phys. Lett. B436 (1998) 403 (2) Phys. Lett. B447 (1999) 147
(3) CERN EP 2000-044 (subm. to Phys. Lett.)

F_2^γ : Tagging in the L3-Detector



- Tagging done in the Luminosity Monitor or in the Endcaps

F_2^γ : Unfolding

- Problem: The final state is not completely reconstructed due to detector acceptance
  - ➪ $W_{vis} < W_{\gamma\gamma}$ and $x_{vis} < x$
- Using kinematics of the tagged electron and transverse momentum conservation can help...
  - ➪ $W_{vis} \rightarrow W_{rec}$
- ... BUT unfolding is still needed!
  - ➪ $W_{rec} \rightarrow W_{\gamma\gamma}$
  - ➪ Dependencies of the used Monte Carlos



F_2^γ : MC - Overview

Used Monte Carlo-Programs

- JAMVG – Vermaseren
- TWOGAM (v1.71) – Nova, Olshevski and Todorov
- PHOJET (v1.05c) – Engel and Ranft
- HERWIG (v5.9) – Marchesini et al.

Parametrisations

- GRV-HO, Glück, Reya and Vogt (1992)
- SaS-1d, Schuler and Sjöstrand (1995, 1996)
- LAC, Levy, Abramovicz and Charchula (1991)

- All Monte Carlos contain different physical models
  - Unfolding depends on the used MC
- This is the main systematic error

F_2^γ : Modeling Dependence



- Unfolding Matrix is specific for the used Monte Carlo
- L3 and OPAL agree well IF the same model is used
- At low x (high mass) the use of different MC's leads to differences in the unfolded values of F_2^γ of 10 to 30 %

F_2^γ : MC Comparison at Low Q^2

Transverse Momentum



- $1.2 < Q^2 < 9$ GeV^2
- PHOJET and TWOGAM describe well the DATA
- HERWIG will not be used

F_2^γ : MC Comparison at Low Q^2

Energy Flow



- ❑ Again PHOJET and TWOGAM describe well the DATA
- ❑ HERWIG agrees better at high x