Standard Model Fits

André Krüger
on behalf of the LEP collaborations
DESY Zeuthen, L3 collaboration

Les Arcs, May 30, 2002

XXXVIIth Rencontres de Moriond

QCD and High Energy Hadronic Interactions

- Introduction
- Measurements
- Fit Results
- Summary

Introduction

Precision electroweak measurements test the Standard Model at the level of its quantum corrections

Modify tree-level relations:

$$G_F = \frac{\pi}{\sqrt{2}} \frac{\alpha(0)}{m_W^2 \sin^2\theta_W} \frac{1}{1-\Delta r}$$

$$\Delta r = \Delta\alpha + \Delta r_W$$

$\Delta\alpha$ absorbed into running of α:

$$\frac{1}{\alpha(0)} \rightarrow \frac{1}{\alpha(m_Z^2)} = \frac{1-\Delta\alpha}{\alpha(0)}$$

Leading electroweak contributions:

$$\Delta r^t \propto m_t^2 \qquad \Delta r^H \propto \log m_H$$









Introduction

$$\Delta\alpha = \Delta\alpha_{\mathrm{lep}} + \Delta\alpha_{\mathrm{t}} + \Delta\alpha^{(5)}_{\mathrm{had}}$$

$\Delta\alpha_{\mathrm{lep}}$: leptonic contribution $\longrightarrow$ precisely calculable
$\Delta\alpha_{\mathrm{t}}$: top quark contribution $\longrightarrow$ small due to high top mass
$\Delta\alpha^{(5)}_{\mathrm{had}}$: light quarks involving QCD

obtain $\Delta\alpha^{(5)}_{\mathrm{had}}$ from dispersion integral:

$$\Delta\alpha^{(5)}_{\mathrm{had}} \propto \mathcal{R}\mathrm{e} \int_{4m_\pi^2}^{\infty} \frac{R(s')}{s'(s'-s)} ds'$$

$$R(s') = \frac{\sigma(e^+e^- \longrightarrow \mathrm{hadrons})}{\sigma(e^+e^- \longrightarrow \mu^+\mu^-)}$$

needs data below 5 GeV and perturbative QCD



Burkhardt/Pietrzyk 2001:

$$\Delta\alpha^{(5)}_{\mathrm{had}} = 0.02761 \pm 0.00036$$

Measurements and Input Parameters for the Fits

LEP I:

- m_Z, Γ_Z, σ^0_{had}, $R_l{}^0$, $A^{0,l}_{FB}$
- $P_\tau(\cos\theta) \rightarrow \mathcal{A}_e$, $\mathcal{A}_\tau$
- R^0_b, R^0_c, $A^{0,b}_{FB}$, $A^{0,c}_{FB}$
- $Q_{FB} \rightarrow \sin^2\theta^{lept}_{eff}$

SLD:

- L-R asymmetry $\rightarrow \mathcal{A}_l$, $\mathcal{A}_b$, $\mathcal{A}_c$

$p\bar{p}$/LEP II:

- m_t, m_W

νN:

- $\rightarrow \sin^2\theta_W$

Theory

$$R^0_l = \Gamma_{had}/\Gamma_l$$

$$R^0_q = \Gamma_q/\Gamma_{had}$$

$$P_\tau(\cos\theta) = -\frac{\mathcal{A}_\tau(1+\cos^2\theta)+2\mathcal{A}_e\cos\theta}{1+\cos^2\theta+2\mathcal{A}_\tau\mathcal{A}_e\cos\theta}$$

$$A^{0,f}_{FB} = \frac{3}{4}\mathcal{A}_e\mathcal{A}_f$$

$$\mathcal{A}_f = \frac{2g^f_V g^f_A}{(g^f_V)^2+(g^f_A)^2}$$

$$g^l_V/g^l_A = 1-4\sin^2\theta^{lept}_{eff}$$

$$\sin^2\theta_W = 1-\frac{m_W{}^2}{m_Z{}^2}$$

Lepton Universality

Final Z lineshape from LEP

$\mathbf{m_Z}$	$\mathbf{91187.5 \pm 2.1\ GeV}$
$\mathbf{\Gamma_Z}$	$\mathbf{2495.2 \pm 2.3\ GeV}$
$\boldsymbol{\sigma^0_{had}}$	$\boldsymbol{41.540 \pm 0.037 nb}$
$\mathbf{R^0_l}$	$\mathbf{20.767 \pm 0.025}$
$\boldsymbol{A^{0,l}_{FB}}$	$\mathbf{0.0171 \pm 0.0010}$



$+\mathbf{P_\tau(\cos\theta)}$ **final:**

$$\mathcal{A}_l^{\mathbf{LEP}} = \mathbf{0.1482 \pm 0.0026}$$

SLD final:

$$\mathcal{A}_l^{\mathbf{SLD}} = \mathbf{0.1513 \pm 0.0021}$$

low $\mathbf{m_H}$ preferred

Measurements: Heavy Flavour

Final results from Z peak

LEP I	R_b^0	0.21646 ± 0.00065
	R_c^0	0.1719 ± 0.0031
	$A_{FB}^{0,b}$	0.0994 ± 0.0017
	$A_{FB}^{0,c}$	0.0685 ± 0.0034
SLD	$\mathcal{A}_b$	0.922 ± 0.020
LEP+SLD	$\mathcal{A}_l$	0.1501 ± 0.0016



Measurements: Heavy Flavour



favours high m_H

Measurements: $\sin^2\theta^{\rm lept}_{\rm eff}$



Leptons vs. Hadrons: 3σ difference

Determination of SM Parameters

LEP I:

- m_Z, Γ_Z, σ^0_{had}, $R_l{}^0$, $A^{0,l}_{FB}$
- $P_\tau(\cos\theta) \rightarrow \mathcal{A}_e$, $\mathcal{A}_\tau$
- R^0_b, R^0_c, $A^{0,b}_{FB}$, $A^{0,c}_{FB}$
- $Q_{FB} \rightarrow \sin^2\theta^{lept}_{eff}$

SLD:

- L-R asymmetry $\rightarrow \mathcal{A}_l$, $\mathcal{A}_b$, $\mathcal{A}_c$

$p\bar{p}$/LEP II:

- m_t, m_W

νN:

- $\rightarrow \sin^2\theta_W$

Theory

$$R^0_l = \Gamma_{had}/\Gamma_l$$

$$R^0_q = \Gamma_q/\Gamma_{had}$$

$$P_\tau(\cos\theta) = -\frac{\mathcal{A}_\tau(1+\cos^2\theta)+2\mathcal{A}_e\cos\theta}{1+\cos^2\theta+2\mathcal{A}_\tau\mathcal{A}_e\cos\theta}$$

$$A^{0,f}_{FB} = \frac{3}{4}\mathcal{A}_e\mathcal{A}_f$$

$$\mathcal{A}_f = \frac{2g^f_V g^f_A}{(g^f_V)^2+(g^f_A)^2}$$

$$g^l_V/g^l_A = 1-4\sin^2\theta^{lept}_{eff}$$

$$\sin^2\theta_W = 1-\frac{m_W{}^2}{m_Z{}^2}$$

χ^2 - fit

m_Z m_t m_H $\alpha(m_Z^2)$ $\alpha_S(m_Z^2)$

Direct Measurement m_W $\leftrightarrow$ SM Prediction



indirect and direct measurement consistent within 1.7σ

Fit Results: m_t and m_W



favours low m_H

Results: m_t, m_W and m_H



prefer low m_H

Fit Results: m_H

$$m_H = 85^{+54}_{-34}\ \text{GeV}$$



Direct searches $m_H > 114.1$ GeV
$m_H < 196$ GeV at 95% C.L.

Fit Results: Global

Winter 2002

	Measurement	Pull
$\Delta\alpha^{(5)}_{had}(m_Z)$	0.02761 ± 0.00036	-.27
m_Z [GeV]	91.1875 ± 0.0021	.01
Γ_Z [GeV]	2.4952 ± 0.0023	-.42
σ^0_{had} [nb]	41.540 ± 0.037	1.63
R_l	20.767 ± 0.025	1.05
$A^{0,l}_{fb}$	0.01714 ± 0.00095	.70
$A_l(P_\tau)$	0.1465 ± 0.0033	-.53
R_b	0.21646 ± 0.00065	1.06
R_c	0.1719 ± 0.0031	-.11
$A^{0,b}_{fb}$	0.0994 ± 0.0017	-2.64
$A^{0,c}_{fb}$	0.0707 ± 0.0034	-1.05
A_b	0.922 ± 0.020	-.64
A_c	0.670 ± 0.026	.06
A_l(SLD)	0.1513 ± 0.0021	1.50
$\sin^2\theta^{lept}_{eff}(Q_{fb})$	0.2324 ± 0.0012	.86
m_W [GeV]	80.451 ± 0.033	1.73
Γ_W [GeV]	2.134 ± 0.069	.59
m_t [GeV]	174.3 ± 5.1	-.08
$\sin^2\theta_W(\nu N)$	0.2277 ± 0.0016	3.00
Q_W(Cs)	-72.39 ± 0.59	.84



$$\text{Pull} = \frac{\mathcal{O}^{\text{meas}} - \mathcal{O}^{\text{fit}}}{\sigma\mathcal{O}^{\text{meas}}}$$

Summary

- the Standard Model describes the data well
- most measurements favour a low Higgs mass (but no Higgs found)
- need precise measurement of $\Delta\alpha^{(5)}_{\mathrm{had}}$ and $\mathrm{m_t}$
- $\sin^2\theta_{\mathrm{W}}$ from NuTeV deviates by 3.0σ

Thanks to the LEPEWWG for the numbers and plots