

Neutral Triple and Quartic Gauge Couplings at LEP

Radoslaw Ofierzynski

ETH Zürich

for the LEP Experiments

7th Topical Seminar on the Legacy of LEP and SLC


October 8 2001

- Neutral Triple Gauge Couplings
 - Theoretical Framework
 - Processes at LEP
 - Results

- Neutral Quartic Gauge Couplings
 - Theoretical Framework
 - Processes at LEP
 - Results

- Summary

Neutral Triple Gauge Couplings





Couplings vanish at tree level in SM.


One-loop level: couplings are of order $\mathcal{O}(10^{-4})$,
below experimental sensitivity.

New Physics \Rightarrow enlargement of neutral TGC's.

Standard Model



Theoretical Framework (neutral TGC)



$$= ie\Gamma_{V_1V_2V_3}^{\alpha,\beta,\mu}(q_1, q_2, P)$$

Vertex parametrisation:
(Lorentz, $U(1)_{em}$ gauge invariance + Bose symmetry)

$$\Gamma_{ZZV}^{\alpha\beta\mu}(q_1, q_2, P) = \frac{i(P^2 - m_V^2)}{m_Z^2} [f_4^V (P^\alpha g^{\mu\beta} + P^\beta g^{\mu\alpha}) - f_5^V \epsilon^{\mu\alpha\beta\rho} (q_1 - q_2)_\rho]$$

$$\Gamma_{Z\gamma V}^{\alpha\beta\mu}(q_1, q_2, P) = \frac{i(P^2 - m_V^2)}{m_Z^2} \left\{ h_1^V (q_2^\mu g^{\alpha\beta} - q_2^\alpha g^{\mu\beta}) + \frac{h_2^V}{m_Z^2} P^\alpha [(P q_2) g^{\mu\beta} - q_2^\mu P^\beta] \right. \\ \left. - h_3^V \epsilon^{\mu\alpha\beta\rho} q_{2\rho} - \frac{h_4^V}{m_Z^2} P^\alpha \epsilon^{\mu\beta\rho\sigma} P_\rho q_{2\sigma} \right\}.$$

(Gounaris et al, hep-ph/0003143,
Hagiwara et al, Nucl.Phys. B282, 253)


- f_4, h_1, h_2 : CP-violating,
amplitudes do NOT interfere with SM
- f_5, h_3, h_4 : CP-conserving,
amplitudes interfere with SM

For nonvanishing contribution: V_1 off-shell.

Neutral Triple Gauge Couplings at LEP


can be tested in $Z\gamma$ and ZZ production.

Experimental NTGC signatures:



test h_i^V

SM background:
2 fermion processes
with ISR



test f_i^V


SM background:
other 4 fermion
processes

$$e^+e^- \rightarrow Z\gamma \text{ (} Z\gamma Z^*, Z\gamma\gamma^* \text{ couplings)}$$

Main effects of anomalous NTGC:

- increase of total cross-section,
- modification of differential spectrum of photon (mainly at large polar angles),
- enhancement of production of longitudinally polarised Z.


DELPHI / U.Baur





$e^+e^- \rightarrow Z\gamma$ ($Z\gamma Z^*$, $Z\gamma\gamma^*$ couplings)

DELPHI Preliminary:

Decay angle of Z in its rest frame in hadronic $Z\gamma$ events




ALEPH Preliminary:






$$e^+e^- \rightarrow ZZ \text{ (} ZZZ^*, ZZ\gamma^* \text{ couplings)}$$

Main effects of anomalous NTGC:

- change in observed total cross-section
- modification of angular distribution of Z
- change in average polarisation of Z



$e^+e^- \rightarrow ZZ$ (ZZZ^* , $ZZ\gamma^*$ couplings)



NTGC Results: $Z\gamma$ Process


ALEPH, DELPHI, L3 use data up to 208GeV
(OPAL only 189GeV)

- ALEPH: (605pb^{-1})
 $q\bar{q}\gamma$: fit using cross-section
 + angular information
- DELPHI: (638pb^{-1})
 $q\bar{q}\gamma$: fit of $d\sigma/d\cos\alpha^*$
 $\nu\bar{\nu}\gamma$: fit cross-section
- L3: (620pb^{-1})
 Optimal Observables:
 $q\bar{q}\gamma$: $(E_\gamma, \theta_\gamma, \phi_\gamma, \theta_f^Z, \phi_f^Z)$
 $\nu\bar{\nu}\gamma$: $(E_\gamma, \theta_\gamma, \phi_\gamma)$
- OPAL: (176pb^{-1})
 $q\bar{q}\gamma$: fit cross section $(E, \cos\theta_\gamma, \cos\alpha)$
 $\nu\bar{\nu}\gamma$: fit cross section $(E, \cos\theta_\gamma)$

Results on h_i^γ ($Z\gamma\gamma^*$ coupling)

Preliminary


LEP **ALEPH+DELPHI+ L3+OPAL**



LEP combination (95% CL limits)


h_1^γ	h_2^γ	h_3^γ	h_4^γ
-0.056	-0.045	-0.049	-0.002
+0.055	+0.025	-0.008	+0.034

Results on h_i^γ (2-dim.)



95% CL

h_1^γ	h_2^γ
-0.16	-0.11
+0.05	+0.02




95% CL

h_3^γ	h_4^γ
-0.08	-0.04
+0.14	+0.11

Results on h_i^Z ($Z\gamma Z^*$ coupling)

Preliminary


LEP **ALEPH+DELPHI+ L3+OPAL**



LEP combination (95% CL limits)


h_1^Z	h_2^Z	h_3^Z	h_4^Z
-0.13	-0.078	-0.20	-0.05
+0.13	+0.071	+0.07	+0.12

Results on h_i^Z (2-dim.)



95% CL

h_1^Z	h_2^Z
-0.35	-0.21
+0.28	+0.17



95% CL

h_3^Z	h_4^Z
-0.37	-0.19
+0.29	+0.21


NTGC Results: ZZ Process

All experiments use data up to 208GeV.

All visible channels used.

- ALEPH: (660pb⁻¹)
fit with cross-section and $\cos\theta_Z$.
- DELPHI: (638pb⁻¹)
fit cross-section.
- L3: (620pb⁻¹)
fit most significant variables,
impact of anomalous couplings from
extension of Excalibur generator.
- OPAL: (609pb⁻¹)
Optimal Observables
+ fit of cross-section


Results on $f_i^{\gamma,Z}$ ($ZZ\gamma^*$, ZZZ^* couplings)

Preliminary
LEP **ALEPH+DELPHI+ L3+OPAL**


LEP combination (95% CL limits)


f_4^γ	f_4^Z	f_5^γ	f_5^Z
-0.17	-0.31	-0.36	-0.36
+0.19	+0.28	+0.40	+0.39

Results on $f_i^{\gamma, Z}$ (2-dim.)



95% CL


f_4^γ	f_4^Z
-0.17	-0.30
+0.19	+0.28



95% CL

f_5^γ	f_5^Z
-0.34	-0.36
+0.38	+0.38


Neutral Quartic Gauge Couplings



Window on Spontaneous Symmetry Breaking.
 New Physics effects due to heavy new particle exchange.

Standard Model

neutral QGC



$$e^+e^- \rightarrow Z\gamma\gamma \rightarrow q\bar{q}\gamma\gamma$$

Theoretical Framework (neutral QGC)

Parametrisation with effective Lagrangian:
(C, P, U(1)_{em}, SU(2)_c symmetries)

$$\mathcal{L}^0 = -\frac{\pi\alpha}{4} \frac{a_0}{\Lambda^2} F_{\mu\nu} F^{\mu\nu} Z^\rho \cdot Z_\rho$$

$$\mathcal{L}^c = -\frac{\pi\alpha}{4} \frac{a_c}{\Lambda^2} F_{\mu\rho} F^{\mu\sigma} Z^\rho \cdot Z_\sigma$$

[Belanger et al., Phys.Lett. B288 (1992)]

α : fine structure constant,

$F_{\mu\nu}$: field strength tensor,

Z^ρ : physical field of Z boson.


Strength of QGC's described by $\frac{a_0}{\Lambda^2}$ and $\frac{a_c}{\Lambda^2}$.
Scale of new physics represented by Λ .

Standard Model: $a_0 = a_c = 0$.

$e^+e^- \rightarrow Z\gamma\gamma$ ($ZZ\gamma\gamma$ coupling)


Main effects of anomalous NQGC:

- deviations in total cross-section,
- less energetic photon gets harder.



$e^+e^- \rightarrow Z\gamma\gamma$ ($ZZ\gamma\gamma$ coupling)

OPAL Preliminary $\sqrt{s} = 130-208$ GeV



(dotted lines: $a_0/\Lambda^2 = 0.015 \text{ GeV}^{-2}$)

NQGC Results: $Z\gamma\gamma$ Process

Problem: new theoretical program (WRAP generator) differs from old one (Stirling&Werthenbach)

Results from July 2001:

(using EEZGG generator)

- L3: (130 – 202GeV, 500 pb⁻¹)
fit to transverse momentum $Pt_{\gamma 2}$
- OPAL: (130 – 208GeV, 580 pb⁻¹)
fit to distribution of
[$E_{\gamma 2}, \max(|\cos \theta_{\gamma 1}|, |\cos \theta_{\gamma 2}|)$]

95% CL limits:

	a_0/Λ^2 (GeV ⁻²)	a_c/Λ^2 (GeV ⁻²)
L3	[-0.008, +0.005]	[-0.007, +0.011]
OPAL	[-0.006, +0.008]	[-0.008, +0.0012]

But: limits are expected to change.

Summary

Neutral Triple Gauge Couplings:

- analyses use $Z\gamma$ and ZZ processes:
- no evidence for neutral TGC's,
- 95% CL limits on $h_i^{\gamma,Z}$, $f_i^{\gamma,Z}$ obtained (LEP combination) of order of 10^{-1} .

Neutral Quartic Gauge Couplings:

- analyses use $Z\gamma\gamma$ process,
- no evidence for neutral QGC's,
- ongoing effort to repeat measurements using WRAP generator.