

Sample	Parameter(s)	Precision
WS/RS $K\pi$	(x'_D, y'_D)	$\mathcal{O}[(10^{-5}, 10^{-4})]$
WS/RS $K\mu\nu$	r_M	$\mathcal{O}(5 \times 10^{-7})$
WS/RS $K\mu\nu$	$ p/q _D$	$\mathcal{O}(1\%)$
$D^{*+} \rightarrow D^0\pi^+ ; D^0 \rightarrow K^-K^+, \pi^-\pi^+$	$\Delta\mathcal{A}_{CP}$	0.015%
$D^{*+} \rightarrow D^0\pi^+ ; D^0 \rightarrow K^-K^+$	\mathcal{A}_{CP}	0.010%
$D^{*+} \rightarrow D^0\pi^+ ; D^0 \rightarrow \pi^-\pi^+$	\mathcal{A}_{CP}	0.015%
$D^{*+} \rightarrow D^0\pi^+ ; D^0 \rightarrow K_S^0\pi^-\pi^+$	(x_D, y_D)	(0.015%, 0.010%)
$D^{*+} \rightarrow D^0\pi^+ ; D^0 \rightarrow K^-K^+, (\pi^-\pi^+)$	y_{CP}	0.004% (0.008%)
$D^{*+} \rightarrow D^0\pi^+ ; D^0 \rightarrow K^-K^+, (\pi^-\pi^+)$	A_Γ	0.004% (0.008%)
$D^{*+} \rightarrow D^0\pi^+ ; D^0 \rightarrow K^-K^+\pi^-\pi^+$	\mathcal{A}_T	2.5×10^{-4}