

Decay	Mode	$j_1$	$j_2$	Allowed values of $h$	Number of amplitudes
$B_s^0 \rightarrow (K^+ \pi^-)_0^* (K^- \pi^+)_0^*$	scalar-scalar	0	0	0	1
$B_s^0 \rightarrow (K^+ \pi^-)_0^* \bar{K}^* (892)^0$	scalar-vector	0	1	0	1
$B_s^0 \rightarrow K^* (892)^0 (K^- \pi^+)_0^*$	vector-scalar	1	0	0	1
$B_s^0 \rightarrow (K^+ \pi^-)_0^* \bar{K}_2^* (1430)^0$	scalar-tensor	0	2	0	1
$B_s^0 \rightarrow K_2^* (1430)^0 (K^- \pi^+)_0^*$	tensor-scalar	2	0	0	1
$B_s^0 \rightarrow K^* (892)^0 \bar{K}^* (892)^0$	vector-vector	1	1	0, $\parallel$ , $\perp$	3
$B_s^0 \rightarrow K^* (892)^0 \bar{K}_2^* (1430)^0$	vector-tensor	1	2	0, $\parallel$ , $\perp$	3
$B_s^0 \rightarrow K_2^* (1430)^0 \bar{K}^* (892)^0$	tensor-vector	2	1	0, $\parallel$ , $\perp$	3
$B_s^0 \rightarrow K_2^* (1430)^0 \bar{K}_2^* (1430)^0$	tensor-tensor	2	2	0, $\parallel_1$ , $\perp_1$ , $\parallel_2$ , $\perp_2$	5