

Source	Uncertainty
τ_h ID	p_T /decay-mode dependent (3%–10%)
τ_h separation from e/μ	3%
$e \rightarrow \tau_h$ ID	η dependent (9%–40%)
$\mu \rightarrow \tau_h$ ID	η dependent (10%–70)%
e ID	2%
μ ID	1%
b jet veto	0–10%
Integrated luminosity	1.6%
Trigger	2% for e/μ , p_T /decay-mode dep. for τ_h [$\mathcal{O}(10\%)$]
$t\bar{t}$ cross section	4.2%
Diboson cross section	5%
Single top quark cross section	5%
Drell-Yan cross section	2%
L1 trigger timing (2016 and 2017)	Event-dependent (0.2%–15%)
$\mathcal{B}(H \rightarrow \tau\tau)$	2.1%
τ_h energy scale	Decay-mode dependent (0.2%–1.2%)
$e \rightarrow \tau_h$ energy scale	Decay-mode dependent (1–7%)
$\mu \rightarrow \tau_h$ energy scale	1%
Electron energy scale	p_T/η dependent ($< 1.25\%$)
Muon energy scale	η dependent 0.4–2.7%
Jet energy scale	p_T/η dependent (0.5%–14%)
Jet energy resolution	η dependent (2%–95%)
p_T^{miss} unclustered energy scale	Event-dependent (0%–20%)
p_T^{miss} recoil corrections	0.3–5.8%
Jet $\rightarrow \tau_h$ misidentification	Event-dependent [$\mathcal{O}(10\%)$]
QCD multijet in the $e\mu$ channel	Event-dependent [$\mathcal{O}(20\%)$]
Embedded yield	4%
$t\bar{t}$ in embedded	10%
Signal theoretical uncertainty	Event-dependent (up to 25%)
Top quark p_T reweighting	p_T dependent (0%–21%)
Drell-Yan p_T and mass reweighting	p_T /mass dependent (0%–11%)