


# INGREDIENTS FOR ACCURATE COLLIDER PHYSICS (1/2)

Gavin Salam, CERN





PSI Summer School Exothiggs,  
Zuoz, August 2016



# The LHC and its Experiments



- ~16.5 mi circumference, ~300 feet underground
- 1232 superconducting twin-bore Dipoles (49 ft, 35 t each)
- Dipole Field Strength 8.4 T (13 kA current), Operating Temperature 1.9K
- Beam intensity 0.5 A ( $2.2 \cdot 10^{-6}$  loss causes quench), 362 MJ stored energy



+ TOTEM, LHCf

## LHC – TWO ROLES – A DISCOVERY MACHINE AND A PRECISION MACHINE

### Today

- 20 fb<sup>-1</sup> at 8 TeV
- 13 fb<sup>-1</sup> at 13 TeV

### Future


- 2018: 100 fb<sup>-1</sup> @ 13 TeV
- 2023: 300 fb<sup>-1</sup> @ 1? TeV
- 2035: 3000 fb<sup>-1</sup> @ 14 TeV

1 fb<sup>-1</sup> = 10<sup>14</sup> collisions


Increase in luminosity brings discovery reach and precision




# The LHC and its Experiments




- ~16.5 mi circumference, ~300 feet underground
- 1232 superconducting twin-bore Dipoles (49 ft, 35 t each)
- Dipole Field Strength 8.4 T (13 kA current), Operating Temperature 1.9K
- Beam intensity 0.5 A ( $2.2 \cdot 10^{-6}$  loss causes quench), 362 MJ stored energy




ATLAS: general purpose



CMS: general purpose



ALICE: heavy-ion physics




LHCb: B-physics

+ TOTEM, LHCf

# LHC – TWO ROLES – A DISCOVERY MACHINE AND A PRECISION MACHINE


## Z' exclusion reach v. lumi




Increase in luminosity brings discovery reach and precision




# The LHC and its Experiments




- ~16.5 mi circumference, ~300 feet underground
- 1232 superconducting twin-bore Dipoles (49 ft, 35 t each)
- Dipole Field Strength 8.4 T (13 kA current), Operating Temperature 1.9K
- Beam intensity 0.5 A ( $2.2 \cdot 10^{-6}$  loss causes quench), 362 MJ stored energy



ATLAS: general purpose



CMS: general purpose



ALICE: heavy-ion physics




LHCb: B-physics

+ TOTEM, LHCf

# LHC – TWO ROLES – A DISCOVERY MACHINE AND A PRECISION MACHINE


## Higgs couplings



Increase in luminosity brings discovery reach and precision



# LONG-TERM HIGGS PRECISION?



Naive extrapolation suggests LHC has long-term potential to do Higgs physics at **1% accuracy**




# THE HIGGS SECTOR

---

The theory is old (1960s-70s).

**But the particle and its theory are unlike anything we've seen in nature.**

- A fundamental scalar  $\phi$ , i.e. spin 0 (all other particles are spin 1 or 1/2)
- A potential  $V(\phi) \sim -\mu^2 (\phi\phi^\dagger) + \lambda(\phi\phi^\dagger)^2$ , which until now was limited to being theorists' "toy model" ( $\phi^4$ )
- "Yukawa" interactions responsible for fermion masses,  $y_i \phi \bar{\psi} \psi$ , with couplings ( $y_i$ ) spanning 5 orders of magnitude






# THE HIGGS SECTOR

The theory is old (1960s-70s).

**But the particle and its theory are unlike anything we've seen in nature.**

- A fundamental scalar  $\phi$ , i.e. spin 0 (all other particles are spin 1 or 1/2)
- A potential  $V(\phi) \sim -\mu^2 (\phi\phi^\dagger) + \lambda(\phi\phi^\dagger)^2$ , which until now was limited to being theorists' "toy model" ( $\phi^4$ )
- "Yukawa" interactions responsible for fermion masses,  $y_i \phi \bar{\psi} \psi$ , with couplings ( $y_i$ ) spanning 5 orders of magnitude



**Higgs sector needs stress-testing**

Is Higgs fundamental or composite?

If fundamental, is it "minimal"?

Is it really  $\phi^4$ ?

Are Yukawa couplings responsible for all fermion masses?




## 3 Signal and background models

The ggF and VBF production modes for  $H \rightarrow WW^*$  are modelled at next-to-leading order (NLO) in the strong coupling  $\alpha_s$  with the PowHEG MC generator [22–25], interfaced with PYTHIA8 [26] (version 8.165) for the parton shower, hadronisation, and underlying event. The CT10 [27] PDF set is used and the parameters of the PYTHIA8 generator controlling the modelling of the parton shower and the underlying event are those corresponding to the AU2 set [28]. The Higgs boson mass set in the generation is 125.0 GeV, which is close to the measured value. The PowHEG ggF model takes into account finite quark masses and a running-width Breit–Wigner distribution that includes electroweak corrections at NLO [29]. To improve the modelling of the Higgs boson  $p_T$  distribution, a reweighting scheme is applied to reproduce the prediction of the next-to-next-to-leading-order (NNLO) and next-to-next-to-leading-logarithm (NNLL) dynamic-scale calculation given by the HRES 2.1 program [30]. Events with  $\geq 2$  jets are further reweighted to reproduce the  $p_T^H$  spectrum predicted by the NLO PowHEG simulation of Higgs boson production in association with two jets ( $H + 2$  jets) [31]. Interference with continuum  $WW$  production [32, 33] has a negligible impact on this analysis due to the transverse-mass selection criteria described in Section 4 and is not included in the signal model.

Jets are reconstructed from topological clusters of calorimeter cells [50–52] using the anti- $k_t$  algorithm with a radius parameter of  $R = 0.4$  [53]. Jet energies are corrected for the effects of calorimeter non-



# ATLAS $H \rightarrow WW^*$ ANALYSIS [1604.02997]



(a)  $N_{\text{jet}} = 0$

That whole paragraph was just for the red part of this distribution (the Higgs signal).

Complexity of modelling each of the backgrounds is comparable



# AIMS OF THESE LECTURES


---

- Give you basic understanding of the “**jargon**” of theoretical collider prediction methods and inputs
- Give you insight into the **power & limitations of different techniques** for making collider predictions




# A proton-proton collision: INITIAL STATE

---




**proton**



**proton**

# A proton-proton collision: FINAL STATE

---



*(actual final-state multiplicity  $\sim$  several hundred hadrons)*



# IT'S MOSTLY QUANTUM CHROMODYNAMICS (QCD)

Quarks — 3 colours:  $\psi_a = \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \end{pmatrix}$

Quark part of Lagrangian:


$$\mathcal{L}_q = \bar{\psi}_a (i\gamma^\mu \partial_\mu \delta_{ab} - g_s \gamma^\mu t_{ab}^C \mathcal{A}_\mu^C - m) \psi_b$$

$SU(3)$  local gauge symmetry  $\leftrightarrow$  8 ( $= 3^2 - 1$ ) generators  $t_{ab}^1 \dots t_{ab}^8$  corresponding to 8 gluons  $\mathcal{A}_\mu^1 \dots \mathcal{A}_\mu^8$ .

A representation is:  $t^A = \frac{1}{2} \lambda^A$ ,

$$\lambda^1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \lambda^2 = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \lambda^3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \lambda^4 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix},$$

$$\lambda^5 = \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix}, \lambda^6 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \lambda^7 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \lambda^8 = \begin{pmatrix} \frac{1}{\sqrt{3}} & 0 & 0 \\ 0 & \frac{1}{\sqrt{3}} & 0 \\ 0 & 0 & \frac{-2}{\sqrt{3}} \end{pmatrix},$$





# IT'S MOSTLY QUANTUM CHROMODYNAMICS (QCD)

Field tensor:  $F_{\mu\nu}^A = \partial_\mu \mathcal{A}_\nu^A - \partial_\nu \mathcal{A}_\mu^A - g_s f_{ABC} \mathcal{A}_\mu^B \mathcal{A}_\nu^C$        $[t^A, t^B] = if_{ABC} t^C$

$f_{ABC}$  are structure constants of  $SU(3)$  (antisymmetric in all indices —  $SU(2)$  equivalent was  $\epsilon^{ABC}$ ). Needed for gauge invariance of gluon part of Lagrangian:

$$\mathcal{L}_G = -\frac{1}{4} F_A^{\mu\nu} F^A_{\mu\nu}$$






# IT'S MOSTLY QUANTUM CHROMODYNAMICS (QCD)

---

The only complete solution uses **lattice QCD**

- put all quark & gluon fields on a 4d lattice  
(NB: imaginary time)
- Figure out most likely configurations  
(Monte Carlo sampling)




*image credit [fdecomite \[flickr\]](#)*

# IT'S MOSTLY QUANTUM CHROMODYNAMICS (QCD)

The only complete solution uses **lattice QCD**

- put all quark & gluon fields on a 4d lattice (NB: imaginary time)
- Figure out most likely configurations (Monte Carlo sampling)

## hadron spectrum from lattice QCD



*Durr et al, arXiv:0906.3599*




# IT'S MOSTLY QUANTUM CHROMODYNAMICS (QCD)

The only complete solution uses **lattice QCD**

- put all quark & gluon fields on a 4d lattice (NB: imaginary time)
- Figure out most likely configurations (Monte Carlo sampling)

## hadron spectrum from lattice QCD



*Durr et al, arXiv:0906.3599*


For LHC reactions, lattice would have to

- Resolve smallest length scales ( $2 \text{ TeV} \sim 10^{-4} \text{ fm}$ )
- Contain whole reaction (pion formed on timescale of  $1 \text{ fm}$ , with boost of  $10000$  — i.e.  $10^4 \text{ fm}$ )

That implies  $10^8$  nodes in each dimension, i.e.  $10^{32}$  nodes — **unrealistic**


# A proton-proton collision: FILLING IN THE PICTURE

---






# A proton-proton collision: FILLING IN THE PICTURE




# A proton-proton collision: SIMPLIFYING IN THE picture

---



- Proton’s dynamics occurs on timescale  $O(1 \text{ fm})$   
Final-state hadron dynamics occurs on timescale  $O(1 \text{ fm})$
- Production of Higgs, Z (and other “**hard processes**”) occurs on timescale  $1/M_H \sim 1/125 \text{ GeV} \sim 0.002 \text{ fm}$



That means we can separate — “**factorise**” — the hard process, i.e. treat it as independent from all the hadronic dynamics




## SHORT-DISTANCE QCD CORRECTIONS ARE PERTURBATIVE

- On timescales  $1/M_H \sim 1/125 \text{ GeV} \sim 0.002 \text{ fm}$  you can take advantage of **asymptotic freedom**
- i.e. you can write results in terms of an expansion in the (*not so*) strong coupling constant  $\alpha_s(125 \text{ GeV}) \sim 0.11$

$$\hat{\sigma} = \hat{\sigma}_0 (1 + c_1 \alpha_s + c_2 \alpha_s^2 + \dots)$$

**LO**  
(Leading Order)




## SHORT-DISTANCE QCD CORRECTIONS ARE PERTURBATIVE

- On timescales  $1/M_H \sim 1/125 \text{ GeV} \sim 0.002 \text{ fm}$  you can take advantage of **asymptotic freedom**
- i.e. you can write results in terms of an expansion in the (*not so*) strong coupling constant  $\alpha_s(125 \text{ GeV}) \sim 0.11$

$$\hat{\sigma} = \hat{\sigma}_0 (1 + \boxed{c_1 \alpha_s} + c_2 \alpha_s^2 + \dots)$$

**NLO**  
(Next-to-Leading Order)




## SHORT-DISTANCE QCD CORRECTIONS ARE PERTURBATIVE

- On timescales  $1/M_H \sim 1/125 \text{ GeV} \sim 0.002 \text{ fm}$  you can take advantage of **asymptotic freedom**
- i.e. you can write results in terms of an expansion in the (*not so*) strong coupling constant  $\alpha_s(125 \text{ GeV}) \sim 0.11$

$$\hat{\sigma} = \hat{\sigma}_0 (1 + c_1 \alpha_s + \boxed{c_2 \alpha_s^2} + \dots)$$


**NNLO**  
(Next-to-next-to-Leading Order)






# THE MASTER EQUATION

$$\sigma(h_1 h_2 \rightarrow ZH + X) = \sum_{n=0}^{\infty} \alpha_s^n \left( \mu_R^2 \right) \sum_{i,j} \int dx_1 dx_2 f_{i/h_1} \left( x_1, \mu_F^2 \right) f_{j/h_2} \left( x_2, \mu_F^2 \right) \times \hat{\sigma}_{ij \rightarrow ZH+X}^{(n)} \left( x_1 x_2 s, \mu_R^2, \mu_F^2 \right) + \mathcal{O} \left( \frac{\Lambda^2}{M_W^4} \right),$$



# THE MASTER EQUATION


$$\sigma(h_1 h_2 \rightarrow ZH + X) = \sum_{n=0}^{\infty} \alpha_s^n(\mu_R^2) \sum_{i,j} \int dx_1 dx_2 f_{i/h_1}(x_1, \mu_F^2) f_{j/h_2}(x_2, \mu_F^2) \times \hat{\sigma}_{ij \rightarrow ZH+X}^{(n)}(x_1 x_2 s, \mu_R^2, \mu_F^2) + \mathcal{O}\left(\frac{\Lambda^2}{M_W^4}\right),$$



*Parton distribution function (PDF): e.g. number of up anti-quarks carrying fraction  $x_2$  of proton's momentum*

# THE MASTER EQUATION

$$\sigma(h_1 h_2 \rightarrow ZH + X) = \sum_{n=0}^{\infty} \alpha_s^n(\mu_R^2) \sum_{i,j} \int dx_1 dx_2 f_{i/h_1}(x_1, \mu_F^2) f_{j/h_2}(x_2, \mu_F^2) \times \hat{\sigma}_{ij \rightarrow ZH+X}^{(n)}(x_1 x_2 s, \mu_R^2, \mu_F^2) + \mathcal{O}\left(\frac{\Lambda^2}{M_W^4}\right),$$




*Parton distribution function (PDF): e.g. number of up quarks carrying fraction  $x_1$  of proton's momentum*



# THE MASTER EQUATION


*Perturbative sum over powers of the strong coupling: typically we use first 2-3 orders*

$$\sigma(h_1 h_2 \rightarrow ZH + X) = \sum_{n=0}^{\infty} \alpha_s^n(\mu_R^2) \sum_{i,j} \int dx_1 dx_2 f_{i/h_1}(x_1, \mu_F^2) f_{j/h_2}(x_2, \mu_F^2) \times \hat{\sigma}_{ij \rightarrow ZH+X}^{(n)}(x_1 x_2 s, \mu_R^2, \mu_F^2) + \mathcal{O}\left(\frac{\Lambda^2}{M_W^4}\right),$$



# THE MASTER EQUATION


$$\sigma(h_1 h_2 \rightarrow ZH + X) = \sum_{n=0}^{\infty} \alpha_s^n \left( \mu_R^2 \right) \sum_{i,j} \int dx_1 dx_2 f_{i/h_1} \left( x_1, \mu_F^2 \right) f_{j/h_2} \left( x_2, \mu_F^2 \right) \times \hat{\sigma}_{ij \rightarrow ZH+X}^{(n)} \left( x_1 x_2 s, \mu_R^2, \mu_F^2 \right) + \mathcal{O} \left( \frac{\Lambda^2}{M_W^4} \right),$$



At each perturbative order  $n$  we have a specific “hard matrix element” (sometimes several for different subprocesses)

# THE MASTER EQUATION

$$\sigma(h_1 h_2 \rightarrow ZH + X) = \sum_{n=0}^{\infty} \alpha_s^n(\mu_R^2) \sum_{i,j} \int dx_1 dx_2 f_{i/h_1}(x_1, \mu_F^2) f_{j/h_2}(x_2, \mu_F^2) \times \hat{\sigma}_{ij \rightarrow ZH+X}^{(n)}(x_1 x_2 s, \mu_R^2, \mu_F^2) + \mathcal{O}\left(\frac{\Lambda^2}{M_W^4}\right),$$



*Additional corrections from non-perturbative effects (higher “twist”, suppressed by powers of QCD scale ( $\Lambda$ ) / hard scale)*



# THE STRONG COUPLING

# RUNNING COUPLING

---

All couplings run (QED, QCD, EW), i.e. they depend on the momentum scale ( $Q^2$ ) of your process.

The QCD coupling,  $\alpha_s(Q^2)$ , runs **fast**:

$$Q^2 \frac{\partial \alpha_s}{\partial Q^2} = \beta(\alpha_s), \quad \beta(\alpha_s) = -\alpha_s^2 (b_0 + b_1 \alpha_s + b_2 \alpha_s^2 + \dots),$$

$$b_0 = \frac{11C_A - 2n_f}{12\pi}, \quad b_1 = \frac{17C_A^2 - 5C_A n_f - 3C_F n_f}{24\pi^2} = \frac{153 - 19n_f}{24\pi^2}$$

Note sign: **Asymptotic Freedom**, due to gluon to self-interaction

2004 Nobel prize: Gross, Politzer & Wilczek


- ▶ At high scales  $Q$ , coupling becomes small
  - ↳ quarks and gluons are almost free, interactions are weak
- ▶ At low scales, coupling becomes strong
  - ↳ quarks and gluons interact strongly — confined into hadrons
  - Perturbation theory fails.

# THE STRONG COUPLING V. SCALE

$$\text{Solve } Q^2 \frac{\partial \alpha_s}{\partial Q^2} = -b_0 \alpha_s^2 \Rightarrow \alpha_s(Q^2) = \frac{\alpha_s(Q_0^2)}{1 + b_0 \alpha_s(Q_0^2) \ln \frac{Q^2}{Q_0^2}} = \frac{1}{b_0 \ln \frac{Q^2}{\Lambda^2}}$$

$\Lambda \simeq 0.2 \text{ GeV}$  (aka  $\Lambda_{QCD}$ ) is the fundamental scale of QCD, at which coupling blows up.


- ▶  $\Lambda$  sets the scale for hadron masses (NB:  $\Lambda$  not unambiguously defined wrt higher orders)
- ▶ Perturbative calculations valid for scales  $Q \gg \Lambda$ .



**PDG World Average:  $\alpha_s(M_Z) = 0.1181 \pm 0.0011$  (0.9%)**

# STRONG-COUPLING DETERMINATIONS

*Bethke, Dissertori & GPS in PDG '16*




- Most consistent set of independent determinations is from lattice
- Two best determinations are from same group (HPQCD, 1004.4285, 1408.4169)  
 $\alpha_s(M_Z) = 0.1183 \pm 0.0007$  (0.6%)  
 [heavy-quark correlators]  
 $\alpha_s(M_Z) = 0.1183 \pm 0.0007$  (0.6%)  
 [Wilson loops]
- Many determinations quote small uncertainties ( $\approx 1\%$ ). All are disputed!
- Some determinations quote anomalously small central values ( $\sim 0.113$  v. world avg. of  $0.1181 \pm 0.0011$ ). Also disputed



# **PARTON DISTRIBUTION FUNCTIONS (PDFs)**

# DEEP INELASTIC SCATTERING

Hadron-hadron is complex because of two incoming partons — so start with simpler Deep Inelastic Scattering (DIS).



Kinematic relations:

$$x = \frac{Q^2}{2p \cdot q}; \quad y = \frac{p \cdot q}{p \cdot k}; \quad Q^2 = xys$$


$$\sqrt{s} = \text{c.o.m. energy}$$

- ▶  $Q^2 =$  photon virtuality  $\leftrightarrow$  *transverse resolution* at which it probes proton structure
- ▶  $x =$  *longitudinal momentum fraction* of struck parton in proton
- ▶  $y =$  momentum fraction lost by electron (in proton rest frame)

# DEEP INELASTIC SCATTERING



$$Q^2 = 25030 \text{ GeV}^2; \quad y = 0.56; \quad \mathbf{x=0.50}$$



# DEEP INELASTIC SCATTERING

---

Write DIS X-section to zeroth order in  $\alpha_s$  ('quark parton model'):

$$\frac{d^2\sigma^{em}}{dx dQ^2} \simeq \frac{4\pi\alpha^2}{xQ^4} \left( \frac{1 + (1-y)^2}{2} F_2^{em} + \mathcal{O}(\alpha_s) \right)$$

$\propto F_2^{em}$  [structure function]

$$F_2 = x(e_u^2 u(x) + e_d^2 d(x)) = x \left( \frac{4}{9} u(x) + \frac{1}{9} d(x) \right)$$

[ $u(x)$ ,  $d(x)$ ): parton distribution functions (PDF)]

NB:

- ▶ use perturbative language for interactions of up and down quarks
- ▶ but distributions themselves have a *non-perturbative* origin.




# PARTON DISTRIBUTION AND DGLAP

- Write up-quark distribution in proton as

$$u(x, \mu_F^2)$$


- $\mu_F$  is the **factorisation scale** — a bit like the renormalisation scale ( $\mu_R$ ) for the running coupling.
- As you vary the factorisation scale, the parton distributions evolve with a renormalisation-group type equation



Dokshitzer-Gribov-Lipatov-Altarelli-Parisi (DGLAP) equations

# DGLAP EQUATION

*take derivative* wrt factorization scale  $\mu^2$



$$\frac{dq(x, \mu^2)}{d \ln \mu^2} = \frac{\alpha_s}{2\pi} \int_x^1 dz p_{qq}(z) \frac{q(x/z, \mu^2)}{z} - \frac{\alpha_s}{2\pi} \int_0^1 dz p_{qq}(z) q(x, \mu^2)$$

$p_{qq}$  is real  $q \leftarrow q$  splitting kernel:  $p_{qq}(z) = C_F \frac{1+z^2}{1-z}$

# DGLAP EQUATION

---

Awkward to write real and virtual parts separately. Use more compact notation:

$$\frac{dq(x, \mu^2)}{d \ln \mu^2} = \frac{\alpha_s}{2\pi} \underbrace{\int_x^1 dz P_{qq}(z) \frac{q(x/z, \mu^2)}{z}}_{P_{qq} \otimes q}, \quad P_{qq} = C_F \left( \frac{1+z^2}{1-z} \right)_+$$

This involves the *plus prescription*:

$$\int_0^1 dz [g(z)]_+ f(z) = \int_0^1 dz g(z) f(z) - \int_0^1 dz g(z) f(1)$$

$z = 1$  divergences of  $g(z)$  cancelled if  $f(z)$  sufficiently smooth at  $z = 1$

# DGLAP EQUATION

Proton contains both quarks and gluons — so DGLAP is a *matrix in flavour space*:

$$\frac{d}{d \ln Q^2} \begin{pmatrix} q \\ g \end{pmatrix} = \begin{pmatrix} P_{q \leftarrow q} & P_{q \leftarrow g} \\ P_{g \leftarrow q} & P_{g \leftarrow g} \end{pmatrix} \otimes \begin{pmatrix} q \\ g \end{pmatrix}$$

[In general, matrix spanning all flavors, anti-flavors,  $P_{qq'} = 0$  (LO),  $P_{\bar{q}g} = P_{qg}$ ]

Splitting functions are:

$$P_{qg}(z) = T_R [z^2 + (1-z)^2], \quad P_{gq}(z) = C_F \left[ \frac{1 + (1-z)^2}{z} \right],$$

$$P_{gg}(z) = 2C_A \left[ \frac{z}{(1-z)_+} + \frac{1-z}{z} + z(1-z) \right] + \delta(1-z) \frac{(11C_A - 4n_f T_R)}{6}.$$

Have various symmetries / significant properties, e.g.

- ▶  $P_{qg}, P_{gg}$ : *symmetric*  $z \leftrightarrow 1-z$  (except virtuals)
- ▶  $P_{qq}, P_{gg}$ : *diverge* for  $z \rightarrow 1$  soft gluon emission
- ▶  $P_{gg}, P_{gq}$ : *diverge* for  $z \rightarrow 0$  Implies PDFs grow for  $x \rightarrow 0$

2015 EPS HEP prize to Bjorken, Altarelli, Dokshitzer, Lipatov & Parisi



NLO:

$$P_{ps}^{(1)}(x) = 4 C_F n_f \left( \frac{20}{9} \frac{1}{x} - 2 + 6x - 4H_0 + x^2 \left[ \frac{8}{3} H_0 - \frac{56}{9} \right] + (1+x) \left[ 5H_0 - 2H_{0,0} \right] \right)$$

$$P_{qg}^{(1)}(x) = 4 C_A n_f \left( \frac{20}{9} \frac{1}{x} - 2 + 25x - 2p_{qg}(-x)H_{-1,0} - 2p_{qg}(x)H_{1,1} + x^2 \left[ \frac{44}{3} H_0 - \frac{218}{9} \right] \right. \\ \left. + 4(1-x) \left[ H_{0,0} - 2H_0 + xH_1 \right] - 4\zeta_2 x - 6H_{0,0} + 9H_0 \right) + 4 C_F n_f \left( 2p_{qg}(x) \left[ H_{1,0} + H_{1,1} + H_2 \right. \right. \\ \left. \left. - \zeta_2 \right] + 4x^2 \left[ H_0 + H_{0,0} + \frac{5}{2} \right] + 2(1-x) \left[ H_0 + H_{0,0} - 2xH_1 + \frac{29}{4} \right] - \frac{15}{2} - H_{0,0} - \frac{1}{2} H_0 \right)$$

$$P_{gq}^{(1)}(x) = 4 C_A C_F \left( \frac{1}{x} + 2p_{gq}(x) \left[ H_{1,0} + H_{1,1} + H_2 - \frac{11}{6} H_1 \right] - x^2 \left[ \frac{8}{3} H_0 - \frac{44}{9} \right] + 4\zeta_2 - 2 \right. \\ \left. - 7H_0 + 2H_{0,0} - 2H_1 x + (1+x) \left[ 2H_{0,0} - 5H_0 + \frac{37}{9} \right] - 2p_{gq}(-x)H_{-1,0} \right) - 4 C_F n_f \left( \frac{2}{3} x \right. \\ \left. - p_{gq}(x) \left[ \frac{2}{3} H_1 - \frac{10}{9} \right] \right) + 4 C_F^2 \left( p_{gq}(x) \left[ 3H_1 - 2H_{1,1} \right] + (1+x) \left[ H_{0,0} - \frac{7}{2} + \frac{7}{2} H_0 \right] - 3H_{0,0} \right. \\ \left. + 1 - \frac{3}{2} H_0 + 2H_1 x \right)$$


$$P_{gg}^{(1)}(x) = 4 C_A n_f \left( 1 - x - \frac{10}{9} p_{gg}(x) - \frac{13}{9} \left( \frac{1}{x} - x^2 \right) - \frac{2}{3} (1+x)H_0 - \frac{2}{3} \delta(1-x) \right) + 4 C_A^2 \left( 27 \right. \\ \left. + (1+x) \left[ \frac{11}{3} H_0 + 8H_{0,0} - \frac{27}{2} \right] + 2p_{gg}(-x) \left[ H_{0,0} - 2H_{-1,0} - \zeta_2 \right] - \frac{67}{9} \left( \frac{1}{x} - x^2 \right) - 12H_0 \right. \\ \left. - \frac{44}{3} x^2 H_0 + 2p_{gg}(x) \left[ \frac{67}{18} - \zeta_2 + H_{0,0} + 2H_{1,0} + 2H_2 \right] + \delta(1-x) \left[ \frac{8}{3} + 3\zeta_3 \right] \right) + 4 C_F n_f \left( 2H_0 \right. \\ \left. + \frac{2}{3} \frac{1}{x} + \frac{10}{3} x^2 - 12 + (1+x) \left[ 4 - 5H_0 - 2H_{0,0} \right] - \frac{1}{2} \delta(1-x) \right) .$$

$$P_{ab} = \frac{\alpha_s}{2\pi} P^{(0)} + \frac{\alpha_s^2}{16\pi^2} P^{(1)}$$

Curci, Furmanski  
& Petronzio '80



# DGLAP evolution (initial quarks only)




Take example evolution starting with just quarks:

$$\partial_{\ln Q^2} q = P_{q \leftarrow q} \otimes q$$

$$\partial_{\ln Q^2} g = P_{g \leftarrow q} \otimes q$$

- ▶ quark is depleted at large  $x$
- ▶ gluon grows at small  $x$

# DGLAP evolution (initial quarks only)




Take example evolution starting with just quarks:

$$\begin{aligned}\partial_{\ln Q^2} q &= P_{q \leftarrow q} \otimes q \\ \partial_{\ln Q^2} g &= P_{g \leftarrow q} \otimes q\end{aligned}$$

- ▶ quark is depleted at large  $x$
- ▶ gluon grows at small  $x$

# DGLAP evolution (initial quarks only)




Take example evolution starting with just quarks:

$$\begin{aligned}\partial_{\ln Q^2} q &= P_{q \leftarrow q} \otimes q \\ \partial_{\ln Q^2} g &= P_{g \leftarrow q} \otimes q\end{aligned}$$

- ▶ quark is depleted at large  $x$
- ▶ gluon grows at small  $x$



# DGLAP evolution (initial quarks only)




Take example evolution starting with just quarks:

$$\partial_{\ln Q^2} q = P_{q \leftarrow q} \otimes q$$

$$\partial_{\ln Q^2} g = P_{g \leftarrow q} \otimes q$$

- ▶ quark is depleted at large  $x$
- ▶ gluon grows at small  $x$

# DGLAP evolution (initial quarks only)




Take example evolution starting with just quarks:

$$\begin{aligned}\partial_{\ln Q^2} q &= P_{q \leftarrow q} \otimes q \\ \partial_{\ln Q^2} g &= P_{g \leftarrow q} \otimes q\end{aligned}$$

- ▶ quark is depleted at large  $x$
- ▶ gluon grows at small  $x$

# DGLAP evolution (initial quarks only)




Take example evolution starting with just quarks:

$$\begin{aligned}\partial_{\ln Q^2} q &= P_{q \leftarrow q} \otimes q \\ \partial_{\ln Q^2} g &= P_{g \leftarrow q} \otimes q\end{aligned}$$

- ▶ quark is depleted at large  $x$
- ▶ gluon grows at small  $x$

# DGLAP evolution (initial quarks only)




Take example evolution starting with just quarks:

$$\begin{aligned}\partial_{\ln Q^2} q &= P_{q \leftarrow q} \otimes q \\ \partial_{\ln Q^2} g &= P_{g \leftarrow q} \otimes q\end{aligned}$$

- ▶ quark is depleted at large  $x$
- ▶ gluon grows at small  $x$

# DGLAP evolution (initial gluons only)



2nd example: start with just gluons.


$$\partial_{\ln Q^2} q = P_{q \leftarrow g} \otimes g$$

$$\partial_{\ln Q^2} g = P_{g \leftarrow g} \otimes g$$

- ▶ gluon is depleted at large  $x$ .
- ▶ high- $x$  gluon feeds growth of small  $x$  gluon & quark.



# DGLAP evolution (initial gluons only)




2nd example: start with just gluons.

$$\partial_{\ln Q^2} q = P_{q \leftarrow g} \otimes g$$

$$\partial_{\ln Q^2} g = P_{g \leftarrow g} \otimes g$$

- ▶ gluon is depleted at large  $x$ .
- ▶ high- $x$  gluon feeds growth of small  $x$  gluon & quark.

# DGLAP evolution (initial gluons only)




2nd example: start with just gluons.

$$\partial_{\ln Q^2} q = P_{q \leftarrow g} \otimes g$$

$$\partial_{\ln Q^2} g = P_{g \leftarrow g} \otimes g$$

- ▶ gluon is depleted at large  $x$ .
- ▶ high- $x$  gluon feeds growth of small  $x$  gluon & quark.

# DGLAP evolution (initial gluons only)




2nd example: start with just gluons.

$$\partial_{\ln Q^2} q = P_{q \leftarrow g} \otimes g$$

$$\partial_{\ln Q^2} g = P_{g \leftarrow g} \otimes g$$

- ▶ gluon is depleted at large  $x$ .
- ▶ high- $x$  gluon feeds growth of small  $x$  gluon & quark.

# DGLAP evolution (initial gluons only)




2nd example: start with just gluons.

$$\partial_{\ln Q^2} q = P_{q \leftarrow g} \otimes g$$

$$\partial_{\ln Q^2} g = P_{g \leftarrow g} \otimes g$$

- ▶ gluon is depleted at large  $x$ .
- ▶ high- $x$  gluon feeds growth of small  $x$  gluon & quark.

# DGLAP evolution (initial gluons only)




2nd example: start with just gluons.

$$\partial_{\ln Q^2} q = P_{q \leftarrow g} \otimes g$$

$$\partial_{\ln Q^2} g = P_{g \leftarrow g} \otimes g$$

- ▶ gluon is depleted at large  $x$ .
- ▶ high- $x$  gluon feeds growth of small  $x$  gluon & quark.

# DGLAP evolution (initial gluons only)



2nd example: start with just gluons.

$$\begin{aligned}\partial_{\ln Q^2} q &= P_{q \leftarrow g} \otimes g \\ \partial_{\ln Q^2} g &= P_{g \leftarrow g} \otimes g\end{aligned}$$

- ▶ gluon is depleted at large  $x$ .
- ▶ high- $x$  gluon feeds growth of small  $x$  gluon & quark.

**DGLAP evolution:**

- ▶ partons lose momentum and shift towards smaller  $x$
- ▶ high- $x$  partons drive growth of low- $x$  gluon




# determining the gluon

---

*which is critical at hadron colliders (e.g. Higgs dominantly produced by gluon-gluon fusion), but not directly probed in Deep-Inelastic-Scattering*

# Consider DIS data – $F_2(x, Q^2)$ – in a world where the proton just had quarks



Fit quark distributions to  $F_2(x, Q_0^2)$ , at *initial scale*  $Q_0^2 = 12 \text{ GeV}^2$ .


NB:  $Q_0$  often chosen lower

Assume there is no gluon at  $Q_0^2$ :

$$g(x, Q_0^2) = 0$$

Use DGLAP equations to evolve to higher  $Q^2$ ; compare with data.

# Consider DIS data – $F_2(x, Q^2)$ – in a world where the proton just had quarks



Fit quark distributions to  $F_2(x, Q_0^2)$ , at *initial scale*  $Q_0^2 = 12 \text{ GeV}^2$ .


NB:  $Q_0$  often chosen lower

Assume there is no gluon at  $Q_0^2$ :

$$g(x, Q_0^2) = 0$$

Use DGLAP equations to evolve to higher  $Q^2$ ; compare with data.

# Consider DIS data – $F_2(x, Q^2)$ – in a world where the proton just had quarks



Fit quark distributions to  $F_2(x, Q_0^2)$ , at *initial scale*  $Q_0^2 = 12 \text{ GeV}^2$ .


NB:  $Q_0$  often chosen lower

Assume there is no gluon at  $Q_0^2$ :

$$g(x, Q_0^2) = 0$$

Use DGLAP equations to evolve to higher  $Q^2$ ; compare with data.

# Consider DIS data – $F_2(x, Q^2)$ – in a world where the proton just had quarks



Fit quark distributions to  $F_2(x, Q_0^2)$ , at *initial scale*  $Q_0^2 = 12 \text{ GeV}^2$ .


NB:  $Q_0$  often chosen lower

Assume there is no gluon at  $Q_0^2$ :

$$g(x, Q_0^2) = 0$$

Use DGLAP equations to evolve to higher  $Q^2$ ; compare with data.

# Consider DIS data – $F_2(x, Q^2)$ – in a world where the proton just had quarks



Fit quark distributions to  $F_2(x, Q_0^2)$ ,  
at *initial scale*  $Q_0^2 = 12 \text{ GeV}^2$ .

NB:  $Q_0$  often chosen lower


Assume there is no gluon at  $Q_0^2$ :

$$g(x, Q_0^2) = 0$$

Use DGLAP equations to evolve to  
higher  $Q^2$ ; compare with data.



# Consider DIS data – $F_2(x, Q^2)$ – in a world where the proton just had quarks



Fit quark distributions to  $F_2(x, Q_0^2)$ , at *initial scale*  $Q_0^2 = 12 \text{ GeV}^2$ .


NB:  $Q_0$  often chosen lower

Assume there is no gluon at  $Q_0^2$ :

$$g(x, Q_0^2) = 0$$

Use DGLAP equations to evolve to higher  $Q^2$ ; compare with data.

# Consider DIS data – $F_2(x, Q^2)$ – in a world where the proton just had quarks



Fit quark distributions to  $F_2(x, Q_0^2)$ , at *initial scale*  $Q_0^2 = 12 \text{ GeV}^2$ .

NB:  $Q_0$  often chosen lower


Assume there is no gluon at  $Q_0^2$ :

$$g(x, Q_0^2) = 0$$

Use DGLAP equations to evolve to higher  $Q^2$ ; compare with data.

**COMPLETE FAILURE  
to reproduce data evolution**

# Consider DIS data – $F_2(x, Q^2)$ – with specially tuned gluon




If gluon  $\neq 0$ , splitting

$$g \rightarrow q\bar{q}$$

generates extra quarks at large  $Q^2$   $\Rightarrow$  faster rise of  $F_2$

Global PDF fits (**CT, MMHT, NNPDF, etc.**) choose gluon distribution that leads to the correct  $Q^2$  evolution.

# Consider DIS data – $F_2(x, Q^2)$ – with specially tuned gluon




If gluon  $\neq 0$ , splitting

$$g \rightarrow q\bar{q}$$

generates extra quarks at large  $Q^2$   $\Rightarrow$  faster rise of  $F_2$

Global PDF fits (CT, MMHT, NNPDF, etc.) choose gluon distribution that leads to the correct  $Q^2$  evolution.

# Consider DIS data – $F_2(x, Q^2)$ – with specially tuned gluon




If gluon  $\neq 0$ , splitting

$$g \rightarrow q\bar{q}$$

generates extra quarks at large  $Q^2$   $\Rightarrow$  faster rise of  $F_2$

Global PDF fits (CT, MMHT, NNPDF, etc.) choose gluon distribution that leads to the correct  $Q^2$  evolution.

# Consider DIS data – $F_2(x, Q^2)$ – with specially tuned gluon



If gluon  $\neq 0$ , splitting


$$g \rightarrow q\bar{q}$$

generates extra quarks at large  $Q^2$   $\Rightarrow$  faster rise of  $F_2$

Global PDF fits (CT, MMHT, NNPDF, etc.) choose gluon distribution that leads to the correct  $Q^2$  evolution.



# Consider DIS data – $F_2(x, Q^2)$ – with specially tuned gluon




If gluon  $\neq 0$ , splitting

$$g \rightarrow q\bar{q}$$

generates extra quarks at large  $Q^2$   $\Rightarrow$  faster rise of  $F_2$

Global PDF fits (CT, MMHT, NNPDF, etc.) choose gluon distribution that leads to the correct  $Q^2$  evolution.

# Consider DIS data – $F_2(x, Q^2)$ – with specially tuned gluon



If gluon  $\neq 0$ , splitting


$$g \rightarrow q\bar{q}$$

generates extra quarks at large  $Q^2$   $\Rightarrow$  faster rise of  $F_2$

Global PDF fits (CT, MMHT, NNPDF, etc.) choose gluon distribution that leads to the correct  $Q^2$  evolution.

**SUCCESS**

# Resulting gluon distribution, compared to quarks



Resulting gluon distribution is **HUGE!**


Carries **47% of proton's momentum**

(at scale of 100 GeV)

Crucial in order to satisfy momentum sum rule.

Large value of gluon has big impact on phenomenology

# Consider DIS data – $F_2(x, Q^2)$ – with specially tuned gluon




If gluon  $\neq 0$ , splitting

$$g \rightarrow q\bar{q}$$

generates extra quarks at large  $Q^2$   $\Rightarrow$  faster rise of  $F_2$

Global PDF fits (**CT, MMHT, NNPDF, etc.**) choose gluon distribution that leads to the correct  $Q^2$  evolution.

# Consider DIS data – $F_2(x, Q^2)$ – with specially tuned gluon




If gluon  $\neq 0$ , splitting

$$g \rightarrow q\bar{q}$$

generates extra quarks at large  $Q^2$   $\Rightarrow$  faster rise of  $F_2$

Global PDF fits (CT, MMHT, NNPDF, etc.) choose gluon distribution that leads to the correct  $Q^2$  evolution.

# Consider DIS data – $F_2(x, Q^2)$ – with specially tuned gluon




If gluon  $\neq 0$ , splitting

$$g \rightarrow q\bar{q}$$

generates extra quarks at large  $Q^2$   $\Rightarrow$  faster rise of  $F_2$

Global PDF fits (CT, MMHT, NNPDF, etc.) choose gluon distribution that leads to the correct  $Q^2$  evolution.

# Consider DIS data – $F_2(x, Q^2)$ – with specially tuned gluon



If gluon  $\neq 0$ , splitting


$$g \rightarrow q\bar{q}$$

generates extra quarks at large  $Q^2$   $\Rightarrow$  faster rise of  $F_2$

Global PDF fits (CT, MMHT, NNPDF, etc.) choose gluon distribution that leads to the correct  $Q^2$  evolution.



# Consider DIS data – $F_2(x, Q^2)$ – with specially tuned gluon



If gluon  $\neq 0$ , splitting

$$g \rightarrow q\bar{q}$$


generates extra quarks at large  $Q^2$   $\Rightarrow$  faster rise of  $F_2$

Global PDF fits (CT, MMHT, NNPDF, etc.) choose gluon distribution that leads to the correct  $Q^2$  evolution.


**SUCCESS**

# TODAY'S PDF FITS





## NNPDF3.0 NLO dataset







## H1 and ZEUS




## Lepton charge asym. v. CT14 @ D0 & CMS



# THREE GLOBAL PDF FITS: CT14, MMHT2014, NNPDF3.0








cross-section ratios  
(W<sup>+</sup>/W<sup>-</sup>, ttbar/Z)  
show tensions with  
some PDFs

ATLAS-CONF-2015-049



*NB: top-quark mass  
choice affects this plot*

# FINAL REMARKS ON PDFS

---

- In range  $10^{-3} < x < 0.1$ , core PDFs (up, down, gluon) known to  $\sim 1\text{-}2\%$  accuracy
- For many LHC applications, you can use PDF4LHC15 set, which merges CT14, MMHT2014, NNPDF30
- Situation is not full consensus: ABM group claims substantially different gluon distribution

For visualisations of PDFs and related quantities,  
a good place to start is


<http://apfel.mi.infn.it/> (ApfelWeb)

# EXTRA SLIDES



# PDFs: WHAT ROUTE FOR PROGRESS?

- Current status is 2–3% for core “precision” region
- Path to 1% is not clear — e.g.  $Z p_T$ 's strongest constraint is on  $qg$  lumi, which is already best known (why?)
- It'll be interesting to revisit the question once  $t\bar{t}bar$ , incl. jets,  $Z p_T$ , etc. have all been incorporated at NNLO
- **Can expts. get better lumi determination? 0.5%?**



# PDF THEORY UNCERTAINTIES

## Theory Uncertainties

