

Resonance	Fit fraction (%)	Real part	Imaginary part	Magnitude	Phase (radians)
$\bar{K}^*(892)^0$	0.88	0.72	0.33	0.03	0.76
$\bar{K}^*(1410)^0$	1.37	0.15	0.22	0.14	1.09
LASS nonresonant	4.09	0.14	0.18	0.11	0.26
$\bar{K}_0^*(1430)^0$	3.32	0.14	0.08	0.07	0.16
LASS total	4.69	—	—	—	—
$\bar{K}_2^*(1430)^0$	1.06	0.26	0.03	0.05	0.65
$\bar{K}^*(1680)^0$	0.80	0.14	0.20	0.11	2.66
$\bar{K}_0^*(1950)^0$	2.42	0.21	0.23	0.22	1.71
$D_{s2}^*(2573)^-$	1.05	—	—	—	—
$D_{s1}^*(2700)^-$	0.54	0.06	0.13	0.04	0.53
$D_{s1}^*(2860)^-$	3.28	0.24	0.09	0.17	0.52
$D_{s3}^*(2860)^-$	0.42	0.05	0.04	0.03	0.18
Nonresonant	7.64	0.28	0.28	0.19	0.48
D_{sv}^{*-}	4.02	0.18	0.17	0.16	0.43
$D_{s0v}^*(2317)^-$	2.30	0.18	0.09	0.13	0.43
B_v^{*+}	1.83	0.25	0.31	0.13	1.53