

Resonance	Fit fraction (%)	Isobar model coefficients			
		Real part	Imaginary part	Magnitude	Phase
$D_0^*(2400)^0$	8.3 ± 2.6	-0.04 ± 0.07	-0.51 ± 0.07	0.51 ± 0.09	-1.65 ± 0.16
$D_2^*(2460)^0$	31.8 ± 1.5	1.00	0.00	1.00	0.00
$D_1^*(2760)^0$	4.9 ± 1.2	-0.32 ± 0.06	-0.23 ± 0.07	0.39 ± 0.05	-2.53 ± 0.24
S-wave nonresonant	38.0 ± 7.4	0.93 ± 0.09	-0.58 ± 0.08	1.09 ± 0.09	-0.56 ± 0.09
P-wave nonresonant	23.8 ± 5.6	-0.43 ± 0.09	0.75 ± 0.09	0.87 ± 0.09	2.09 ± 0.15
$D_v^*(2007)^0$	7.6 ± 2.3	0.16 ± 0.08	0.46 ± 0.09	0.49 ± 0.07	1.24 ± 0.17
B_v^*	3.6 ± 1.9	-0.07 ± 0.08	0.33 ± 0.07	0.34 ± 0.06	1.78 ± 0.23
Total fit fraction	118.1				