

Resonance	Spin	DP axis	Model	Parameters
$K^*(892)^0$	1	$m^2(K^+\pi^-)$	RBW	$m_0 = 895.81 \pm 0.19 \text{ MeV}$ , $\Gamma_0 = 47.4 \pm 0.6 \text{ MeV}$
$K^*(1410)^0$	1	$m^2(K^+\pi^-)$	RBW	$m_0 = 1414 \pm 15 \text{ MeV}$ , $\Gamma_0 = 232 \pm 21 \text{ MeV}$
$K_0^*(1430)^0$	0	$m^2(K^+\pi^-)$	LASS	Determined from data (see text)
$K_2^*(1430)^0$	2	$m^2(K^+\pi^-)$	RBW	$m_0 = 1432.4 \pm 1.3 \text{ MeV}$ , $\Gamma_0 = 109 \pm 5 \text{ MeV}$
$D_0^*(2400)^-$	0	$m^2(\bar{D}^0\pi^-)$	RBW	Determined from data (see Table ??)
$D_2^*(2460)^-$	2	$m^2(\bar{D}^0\pi^-)$	RBW	
Nonresonant	0	$m^2(\bar{D}^0\pi^-)$	dabba	Fixed (see text)
Nonresonant	1	$m^2(\bar{D}^0\pi^-)$	EFF	Determined from data (see text)