

Ratio	$\bar{D}^{*0} \rightarrow \bar{D}^0 \gamma$	$\bar{D}^{*0} \rightarrow \bar{D}^0 \pi^0$
$\frac{\mathcal{B}(B^0 \rightarrow \bar{D}^{*0} K^+ \pi^-)}{\mathcal{B}(B^0 \rightarrow \bar{D}^{*0} \pi^+ \pi^-)}$	$0.083 \pm 0.005 \pm 0.008$	$0.085 \pm 0.006 \pm 0.012$
$\frac{\mathcal{B}(B_s^0 \rightarrow \bar{D}^{*0} K^- \pi^+)}{\mathcal{B}(B^0 \rightarrow \bar{D}^{*0} \pi^+ \pi^-)}$	$1.19 \pm 0.03 \pm 0.10 \pm 0.04$	$1.10 \pm 0.04 \pm 0.13 \pm 0.03$
$\frac{\mathcal{B}(B^0 \rightarrow \bar{D}^{*0} K^+ \pi^-)}{\mathcal{B}(B_s^0 \rightarrow \bar{D}^{*0} K^- \pi^+)}$	$0.070 \pm 0.004 \pm 0.008 \pm 0.002$	$0.077 \pm 0.006 \pm 0.011 \pm 0.002$